version September 29, 2016: fm
Preprint typeset using LaTeX style emulateapj v. 11/10/09

arXiv:1609.08615v1 [astro-ph.HE] 27 Sep 2016

THE INFRARED–GAMMA-RAY CONNECTION. A *WISE* VIEW OF THE EXTRAGALACTIC GAMMA-RAY SKY

F. Massaro[1,2,3] & R. D'Abrusco[4]
Dipartimento di Fisica, Università degli Studi di Torino, via Pietro Giuria 1, I-10125 Torino, Italy
Istituto Nazionale di Fisica Nucleare, Sezione di Torino, I-10125 Torino, Italy
Istituto Nazionale di Astrofisica - Osservatorio Astrofisico di Torino, via Osservatorio 20, I-10025 Pino Torinese, Italy and
Smithsonian Astrophysical Observatory, 60 Garden Street, Cambridge, MA 02138, USA
version September 29, 2016: fm

ABSTRACT

Using data from the *WISE* all-sky survey we discovered that the non-thermal infrared (IR) emission of blazars, the largest known population of extragalactic γ-ray sources, has peculiar spectral properties. In this work, we confirm and strengthen our previous analyses using the latest available releases of both the *WISE* and the *Fermi* source catalogs. We also show that there is a tight correlation between the mid-IR colors and the γ-ray spectral index of *Fermi* blazars. We name this correlation *the infrared–γ-ray connection*. We discuss how this connection links both the emitted powers and the spectral shapes of particles accelerated in jets arising from blazars over ten decades in energy. Based on this evidence, we argue that *the infrared–γ-ray connection* is stronger than the well known *radio–γ-ray connection*.

Subject headings: galaxies: active - galaxies: BL Lacertae objects: general - radiation mechanisms: non-thermal - gamma rays: general - infrared: general

1. INTRODUCTION

Blazars are one of the most extreme class of radio-loud active galaxies whose emission extends from radio to TeV energies. They generally show extreme variability at all wavelengths and with timescales spanning from weeks to minutes, evidence of superluminal motions, high and variable polarization, flat radio spectra (see e.g. Urry & Padovani 1995), recently observed even below ∼1GHz (i.e. , Massaro et al. 2013a; Massaro et al. 2013d) and a characteristic double bumped spectral energy distribution (SED; see also Massaro et al. 2009, for a recent review).

Since the launch of the *Fermi* satellite (Atwood et al. 2009), blazars have been identified as the dominant class of γ-ray sources, not only extragalactic. Blazars account for about 1/3 of the *Fermi* detected objects (Acero et al. 2015) and likely for a significant fraction of the unidentified/unassociated γ-ray sources (UGSs; Massaro et al. 2012a; Massaro et al. 2013b). Together with star forming regions (e.g. Ackermann et al. 2012a) and radio galaxies (e.g. Di Mauro et al. 2014; Massaro & Ajello 2011b), blazars produce a significant contribution to the extragalactic γ-ray background (Ajello et al. 2012; Ajello et al. 2014; Massaro et al. 2016, and references therein).

At optical frequencies blazars are historically split in two subclasses: BL Lac objects and flat spectrum radio quasars. The former, that will be indicated in this paper as BZBs following the nomenclature introduced by the *Roma-BZCAT* (Massaro et al. 2015b), show featureless spectra and/or with weak absorption lines of equivalent width lower than $5\AA$, while the latter, usually indicated as BZQs, have quasar-like optical spectra (Stickel et al. 1991; Falomo et al. 2014).

Since 2010, the NASA Wide-field Infrared Survey Explorer (*WISE*; Wright et al. 2010) has mapped the sky in the infrared (IR) at 3.4, 4.6, 12, and 22 μm, making possible the investigation of the mid-IR properties of a large, statistically significant, sample of confirmed blazars. We discovered that *Fermi* blazars inhabit a region of the mid-IR color-color diagram, built with the *WISE* magnitudes, well separated from the location of other extragalactic sources (Massaro et al. 2011a; D'Abrusco et al. 2012). This 2-dimensional region in the mid-IR color-color diagram [3.4]-[4.6]-[12] μm was originally indicated as the *WISE Gamma-ray Strip*, and it is the projection of a 3-dimensional volume in the [3.4]-[4.6]-[12]-[22] μm mid-IR color space known as the *WISE locus* of γ-ray blazars (D'Abrusco et al. 2013; D'Abrusco et al. 2014).

These findings led to the development of different procedures to search for γ-ray blazar candidates within the positional uncertainty regions of the *Fermi* UGSs that selected hundreds of IR sources as candidate blazars. Thanks to an extensive optical spectroscopic follow-up campaign (Paggi et al. 2014; Massaro et al. 2014; Massaro et al. 2015a) we confirmed the nature of hundreds of new γ-ray blazars (see also Massaro et al. 2015c; Landoni et al. 2015; Ricci et al. 2015) and assessed the reliability of our association methods. Optical spectroscopy was also used to determine the nature of *Fermi* sources classified as active galaxies of uncertain type (AGUs; Nolan et al. 2012; Ackermann et al. 2011) and/or blazar candidates of uncertain type (BCUs; Acero et al. 2015; Ackermann et al. 2015a).

Here we present an update of the *WISE Gamma-ray Strip* obtained by combining the latest releases of both the *WISE* All-Sky catalog[1] and the Fermi Large Area Telescope Third Source Catalog (3FGL; Acero et al. 2015). Additionally, for the first time, we discuss the link found between the *WISE* mid-IR colors and the γ-ray photon index, comparing it with the radio–γ-ray connection (e.g. Ackermann et al. 2011b).

For our numerical results, we use cgs units unless stated otherwise. Gamma-ray photon index, Γ is defined by the usual convention on the flux density, $N(E) \propto$

[1] http://wise2.ipac.caltech.edu/docs/release/allsky/

$E^{-\Gamma}$, being $N(E)$ the number of γ-ray photons detected per unit of time, area and energy. *WISE* magnitudes are in the Vega system and are not corrected for the Galactic extinction since, as shown in our previous analyses, such correction affects significantly only the magnitude at 3.4μ for sources lying at low Galactic latitudes (see e.g. D'Abrusco et al. 2014). *WISE* bands are indicated as $w1$, $w2$, $w3$ and $w4$ and corresponds to the following nominal wavelengths: 3.4μm, 4.6μm, 12μm and 22μm, respectively.

2. SAMPLE SELECTION

The analysis presented hereby has been carried out using three distinct samples of sources extracted from the *Roma-BZCAT* catalog of confirmed blazars, the 3FGL catalog of *Fermi* sources and catalog of radio sources extracted from the NVSS and SUMSS surveys, respectively. Then, we searched the mid-IR counterparts in the *WISE* all-sky catalog for all sources in each of these samples (the fraction of objects in each sample with a *WISE* association are reported reported in Table 1). It is worth noting that the Two Micron All Sky Survey (2MASS) (Skrutskie et al. 2006) counterparts of the *WISE* sources are automatically reported in the *WISE* catalog.

The first sample includes all BZBs and BZQs listed in the latest version of the *Roma-BZCAT* (i.e., v5.0; Massaro et al. 2015b) and associated to a *Fermi* source belonging to the Third Catalog of Active Galactic Nuclei Detected by the *Fermi* Large Area Telescope (3LAC Ackermann et al. 2015a). This sample is the update of the sample used to build the *WISE Gamma-ray Strip* in (D'Abrusco et al. 2012), and contains 1036 γ-ray blazars: 610 BZBs and 426 BZQs. Adopting the statistical criterion described in our previous analyses we identified as blazar counterpart the closest IR source within $3''.3$ from the position reported in the *Roma-BZCAT* (e.g., D'Abrusco et al. 2014). This criterion was chosen because source positions of the *Roma-BZCAT*, even if often more precise than those reported in the 3LAC, are taken from various catalogs with different positional uncertainties. Given the high detection rate of *Fermi* blazars in the first 3 *WISE* filters we only focus on the analysis in the color diagram built with the 3.4μm, 4.6μm and 12μm magnitudes.

The second sample is composed by BCUs of the 3FGL with a *WISE* counterpart within $10''$. This is the typical radio positional uncertainty used in the 3LAC, usually corresponding to that of the radio NRAO VLA Sky Survey (NVSS Condon et al. 1998) and/or the Sydney University Molonglo Sky Survey (SUMSS; Mauch et al. 2003) used to search for counterparts of *Fermi* sources.

The third sample includes all the radio sources of both the NVSS and the SUMSS catalog lying within the positional uncertainty region of the UGSs at 95% level of confidence and with a *WISE* counterpart detected at least at 3.4μm, 4.6μm and 12μm, within a maximum radius obtained by adding the radio to the mid-IR positional uncertainties at 1σ level of confidence.

3. GAMMA-RAY BLAZARS AT IR WAVELENGTHS

The extremely high detection rate of *Fermi* blazars at mid-IR frequencies (≈99%) at 3.4 and 4.6 μm (see Table 1), strongly supports the use of statistical methods combined with optical spectroscopic campaigns toward the identification of *WISE* selected blazar-like sources that can be responsible for the UGSs (Paggi et al. 2013; Acero et al. 2013).

TABLE 1
NUMBER OF SOURCES DETECTED AT IR FREQUENCIES IN BOTH THE *WISE* AND THE 2MASS CATALOGS AND FOR THE THREE SAMPLES CONSIDERED IN OUR ANALYSIS.

Sample	$w1$	$w2$	$w3$	$w4$	J	H	K	Total
BZB	603	603	595	497	521	523	528	610
BZQ	419	419	419	405	207	206	215	426
BZB ∪ BZQ	1022	1022	1014	902	728	729	743	1036
BCU	569	569	553	433	319	328	325	571
UGS	1546	1465	558	315	580	607	584	1548

Note: *WISE* bands are indicated as $w1$, $w2$, $w3$ and $w4$ while 2MASS magnitudes are J, H, K.

The distribution of the updated sample of *Fermi* blazars in the WISE color-color diagram shown in Fig. 1, confirms our previous results (see e.g. Massaro et al. 2011a) with a 5 times larger sample. The use of *WISE* colors as distinct markers of the non-thermal emission of blazars is clearly supported by these findings.

For comparison, in the color-color diagram generated using the J,H and K magnitudes in the 2MASS catalog (see Fig. 1), although a clear separation between generic IR sources and the region preferentially populated by *Fermi* blazars is visible, no peculiar color pattern is observed for BZBs and BZQs. Moreover, a large fraction of the *Fermi* BZQs considered do not have a 2MASS counterparts (see Table 1), strongly limiting the potential efficiency and completeness of methods for the association of UGS based on 2MASS photometry. The region occupied by the *Fermi* blazars in the 2MASS color-color plot is similar to that previously indicated by Chen et al. (2005) for a different sample of BZBs and BZQs. However, about 30% of the blazars in the Chen et al. (2005) sample show near-IR colors corresponding to a blackbody spectral shape, suggesting a possible contributions from their host galaxies. Such strong contamination is not present at mid-IR wavelengths.

4. THE INFRARED–GAMMA-RAY CONNECTION

To assess the significance of the *infrared–γ-ray connection*, we performed a statistical analysis of the correlation between the γ-ray photon index Γ of the *Fermi* blazars and the *WISE* mid-IR colors, since colors are a surrogate of the spectral index. We found a strong correlation between Γ and both the *WISE* colors [3.4]-[4.6] μm and [4.6]-[12] μm as shown in Fig. 2. This result constitutes the first direct evidence of the existence of the *IR–γ-ray connection*.

In both cases, the correlation is more statistically significant for BZBs rather than for BZQs. The linear correlation coefficient ρ estimated for the Γ vs [3.4]-[4.6] μm correlation is 0.47 for BZBs, with negligible probability to be spurious. On the other hand, for the BZQs, we obtained ρ =0.15 calculated with 419 sources and still with a negligible chance probability, given the large number of objects considered. When using the data for the whole sample of 1022 *Fermi* blazars, the linear correlation coefficient, unsurprisingly, increases to ρ =0.65 thanks to the quite distinct distributions of both BZBs and BZQs



FIG. 1.— Left) Distribution of sources belonging to the *WISE Gamma-ray Strip* (see § 2), in the $w2-w3$ vs $w1-w2$ color-color diagram. BZBs and BZQ are displayed as blue circles and red squares respectively. ∼3000 generic IR sources (grey circles) selected at high Galactic latitudes are also show. The separation between the mid-IR colors of the *Fermi* blazars and that of other sources randomly chosen appears evident. Right) Distribution of the same sample in the near-IR 2MASS $H-K$ vs $J-H$ color-color diagram. The distinction between the region occupied by the *Fermi* blazars and other celestial objects appears clear also in this case but there is no neat separation between BZBs and BZQs.

in the Γ *vs* [3.4]-[4.6] μm plane.

A similar situation is observed in the Γ *vs* [4.6]-[12] μm plane, where the correlation coefficients for the 595 BZBs and the 419 BZQs separately are ρ =0.48 and ρ =0.13 respectively, both with negligible chance probability. The correlation coefficient for the whole *Fermi* blazar sample (1014 sources) is ρ=0.60, lower than the previous one due to the scatter in the plane introduced by the larger uncertainties on the mid-IR magnitudes at 12μm. The slope of the best fit regression lines in both cases, as expected, is driven by the stronger correlations found for the BZBs.

Finally, we did not find any significant trend between the 2MASS J, H and K colors, corrected for Galactic extinction, and the *Fermi* γ-ray spectral index Γ.

5. COMPARISON WITH THE RADIO–GAMMA-RAY CONNECTION

Since the release of the first observations carried out with the *Compton Gamma-ray Observatory*, a link between radio and γ-ray emissions of blazars appeared evident (e.g., Stecker et al. 1993). This well established *radio–γ-ray connection* has mostly driven the searches for counterparts associable with γ-ray sources (Mattox et al. 1997), and has been invoked to justify radio-follow up observations of the positional uncertainty regions for UGSs to date (e.g., Schinzel et al. 2015, and references therein). This connection between the blazar emission at ∼ 1GHz and at GeV energies is due to the correlation between their flux densities and, consequently, their powers.

Thanks to the large increase of the number of blazars detected by *Fermi* it has become possible to study the presence of biases and selection effects that have to be taken into account to confirm the radio–γ-ray connection. Intrinsic variability, redshift dependence, source misidentifications and incorrect associations (see e.g. Mucke et al. 1997), to name a few effects that can mimic a spurious trend, were not found to affect this correlation. Based on this analysis, the existence of the radio–γ-ray connection (see also Mahony et al. 2010, and references therein) was recently confirmed by Ackermann et al. (2011b).

For the sake of comparison, it is worth stressing that the *radio–γ-ray connection* is based on the correlation between the radio and γ-ray flux densities. A similar trend also exists for the mid-IR emission as shown in the lower panels of Fig. 2, which display the 3.4 μm and the 12μm magnitudes vs γ-ray flux density for both confirmed *Fermi* BZBs and BZQs.

Based on the discussion in the previous Section, it appears evident that the *IR–γ-ray connection* is stronger than the radio–γ-ray one, because it is not only a link between flux densities but it also involves a tight connection between spectral shapes of the emitting particles.

6. THE UNCERTAIN GAMMA-RAY SKY

We stress the strength of the *IR–γ-ray connection* by showing that potential counterparts of 3FGL UGSs selected via their WISE Mid-IR colors are in good agreement with the correlation between the [3.4] and [12] WISE colors and the γ-ray Γ spectral index discussed in § 4.

We considered all the *WISE* sources with a radio counterpart that lie within the positional uncertainty regions of the UGSs at 95% level of confidence. By applying the Kernel Density Estimation (KDE) technique (see e.g. D'Abrusco et al. 2009, and reference therein), as in our previous analyses (see e.g., Massaro et al. 2011a; Massaro et al. 2013c; Massaro et al. 2015a), we estimated the probability density function of the IR color distribution for the *Fermi* blazar population and we selected as blazar-like candidates those located within the 95% isodensity contours drawn from the KDE probabil-



FIG. 2.— Upper panels): the Γ *vs* [3.4]-[4.6] μm (left) and the Γ *vs* [4.6]-[12] μm (right) diagrams of the *WISE Gamma-ray Strip* sample. BZBs are shown in blue circles while BZQs in red squares. The correlations between the γ-ray spectral shape and the *WISE* mid-IR colors are highlighted by the regression lines (black) for the whole sample and for BZB and BZQ subclasses separately. Lower panels): distribution of the *WISE Gamma-ray Strip* sources in the wse magnitudes at [3.4] (left) and [12] (right) μm vs γ-ray flux density planes. The distribution on the right shows that 12 μm emission from BZBs is less contaminated by the host galaxy than at 3.4 μm.

ities in the [3.4]-[4.6]-[12] μm diagnostic diagram.

In Fig. 3 we show, as an example, the 8 *WISE* sources with a radio counterpart that lie within the positional uncertainty region at 95% level of confidence of the UGS 3FGL J1216.6-0557. The only *WISE* source selected by the KDE analysis is the closest to the regression line for the *WISE* [3.4]-[4.6] μm vs γ-ray spectral shape distribution of all the *Fermi* blazars (BZBs and BZQs) considered in our analysis (see § 2 and § 4 for more details).

In general, by applying the same technique to all the 987 UGSs listed in the 3FGL to date, we selected 130 blazar-like potential counterparts out of 1548 *WISE* sources with a radio counterpart lying within the 95% positional uncertainty regions. As shown in the Γ *vs* [3.4]-[4.6] μm diagram (Fig. 4, top left), these candidates are clearly located in the region occupied by the *Fermi* blazars, thus following the *IR-γ-ray connection*. It is worthwhile to stress that such consistency it is not granted *a priori* since the *WISE* colors of *Fermi* blazars simply reflect the fact that their mid-IR emission is dominated by non-thermal radiation independently of any γ-ray information. As a consequence, the *IR–gamma-ray connection* cannot be a consequence of the existence of *WISE* γ-ray Strip.

To complete our investigation we also analyzed the 3FGL BCUs. As expected, the mid-IR colors of a large fraction of them (i.e., more than 90%) are similar to the *WISE* colors of confirmed *Fermi* blazars and follow the correlation between Γ and [3.4]-[4.6] μm (4) that we have denominated the *IR–γ-ray connec-*



FIG. 3.— The Γ *vs* [3.4]-[4.6] μm distribution of WISE sources with a radio counterpart within the region of positional uncertainty at 95% level of confidence for the UGS 3FGL J1216.6-0557 (squares). Orange circles in the background are confirmed *Fermi* blazars and the black line shows the regression between their spectral parameters. The arrows indicate the only two *WISE* sources detected at [3.4]-[4.6]-[12] μm for which it was possible to compute the KDE probability. The only *WISE* source selected with the KDE method is the closest to the regression line correspondent to the *IR–γ-ray connection*.

tion. A large fraction of these blazar-like sources are also expected to be BZBs, in agreement with the recent results of several optical spectroscopic campaigns (see e.g. Shaw et al. 2013; Álvarez-Crespo et al. 2016a; Álvarez-Crespo et al. 2016b, for more details) which find that more than 70% of the observed sources are BZBs.

7. SUMMARY AND DISCUSSION

Five years after the discovery that the non-thermal emission of γ-ray blazars can be traced using mid-IR colors obtained from the photometry of the *WISE* all-sky survey, we present an updated analysis based on the latest releases of both the *WISE* and the *Fermi* catalogs. Then, for the first time, we discuss on the existence of a *IR–γ-ray connection* for the *Fermi* blazars that appears to be at least as strong as the well-known radio–γ-ray one.

Our results can be summarized as follows.

1. Using the largest sample of *Fermi* blazars available to date, we confirmed that this extreme class of active galaxies occupied a narrow and well defined region in the mid-IR color-color plane, the so called *WISE Gamma-ray Strip*.

2. Comparing mid-IR *WISE* and near-IR 2MASS diagnostic diagrams, we confirmed that in the latter *Fermi* blazar are distinct from generic 2MASS sources, even though no clear color-color trend appears as in the former. *Fermi* blazars have a low detection rate in the 2MASS catalog, mainly due the 2MASS higher flux limit compared to *WISE* survey. These reasons advice against the use of near-IR colors to search for potential blazar-like counterparts of the UGSs.

3. We describe the statistically significant correlations between the γ-ray photon index and the mid-IR colors for both the whole sample of *Fermi* blazars and the BZBs and BZQs spectral classes separately, the basis of the *IR–γ-ray connection*. This correlation appears "stronger" than the well known radio–γ-ray connection because it involves the spectral shapes of the *Fermi* blazars over $\sim$10 orders of magnitude and not only their flux densities.

4. We argue that the peculiar mid-IR colors of *Fermi* blazars do not depend on their γ-ray photon index. In turn, the *IR–γ-ray connection is* unexpected. We have also highlighted its potential use to implement the search for blazar-like counterparts of the *Fermi* UGSs.

5. We show that a large fraction of the BCUs listed in the 3FGL whose *WISE* counterpart have mid-IR colors consistent with the *WISE Gamma-ray Strip*, have *Fermi* spectral index values consistent with the *IR–γ-ray connection*.

Taking advantage of the overwhelming fraction of *Fermi* blazars detected in the *WISE* all-sky survey (i.e., $\sim$99%) in the first two mid-IR filters, we suggest that a comprehensive investigation of their IR properties at the light of the *IR–γ-ray connection* described in this paper, can represent a powerful tool to reveal the real fraction of *Fermi* blazars hidden within the sample of UGSs.

We thank the anonymous referee for useful comments that led to improvements in the paper. F.M. gratefully acknowledges the financial support of the Programma Giovani Ricercatori – Rita Levi Montalcini – Rientro dei Cervelli (2012) awarded by the Italian Ministry of Education, Universities and Research (MIUR). This research has made use of data obtained from the high-energy Astrophysics Science Archive Research Center (HEASARC) provided by NASA's Goddard Space Flight Center. The NASA/IPAC Extragalactic Database (NED) operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. Part of this work is based on the NVSS (NRAO VLA Sky Survey): The National Radio Astronomy Observatory is operated by Associated Universities, Inc., under contract with the National Science Foundation and on the VLA Low-frequency Sky Survey (VLSS). The Molonglo Observatory site manager, Duncan Campbell-Wilson, and the staff, Jeff Webb, Michael White and John Barry, are responsible for the smooth operation of Molonglo Observatory Synthesis Telescope (MOST) and the day-to-day observing programme of SUMSS. The SUMSS survey is dedicated to Michael Large whose expertise and vision made the project possible. The MOST is operated by the School of Physics with the support of the Australian Research Council and the Science Foundation for Physics within the University of Sydney. This publication makes



FIG. 4.— The [3.4]-[4.6]-[12] μm color-color diagrams (left panels) and the Γ *vs* [3.4]-[4.6] μm diagrams (right panels) for blazar-like sources associated to 3FGL UGSs (upper panels) and 3FGL BCUs (lower panels), respectively. A large fraction of BCUs (i.e., more than 90%) are consistent with the *WISE* colors of confirmed *Fermi* blazars, i.e. *WISE Gamma-ray Strip* (lower left). The dashed ellipse highlights a small number of BCUs whose *WISE* counterparts have mid-IR colors similar to those of normal galaxies and do not follow the correlation between the IR and the gamma-ray spectral shape (lower right). In the upper left panel we show the comparison between all the *WISE* sources lying within the positional uncertainty regions of the 3FGL UGSs with a radio counterpart (magenta diamonds) and the sources of the *WISE Gamma-ray Strip*. The *WISE* sources that are selected according to the KDE method are shown as black circles. These *WISE* sources are mostly consistent with the IR–γ-ray connection (upper right). In all panels *Fermi* blazars are shown as orange circles regardless of their spectral class. Generic mid-IR sources are shown in the background as grey circles in the two panels on the left.

use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. TOPCAT[2] (Taylor 2005) for the preparation and manipulation of the tabular data and the images.

[2] http://www.star.bris.ac.uk/~mbt/topcat/

REFERENCES

Atwood, W. B., Abdo, A. A., Ackermann, M. et al. 2009 ApJ, 697, 1071
Acero, F., Donato, D., Ojha, R. et al. 2013 ApJ, 779, 133
Acero, F., Ackermann, M., Ajello, M. et al. 2015 ApJS, 218, 23

Ackermann, M., Ajello, M., Allafort, A. et al. 2011a ApJ, 743, 171
Ackermann, M., Ajello, M., Allafort, A. et al. 2011b ApJ, 741, 30
Ackermann, M., Ajello, M., Allafort, A. et al. 2012a ApJ, 755, 164
Ackermann, M., Ajello, M., Atwood, W. B. et al. 2015 ApJ, 810, 14
Ajello, M., Shaw, M. S., Romani, R. W. et al. 2012 ApJ, 751, 108
Ajello, M. et al. 2014 ApJ, 780, 73
Álvarez-Crespo, N., Masetti, N., Ricci, F. et al. 2016a AJ in press
Álvarez-Crespo, N., Massaro, F., Milisavljevic, D. et al. 2016b AJ in press
Chen, P. S., Fu, H. W., Gao, Y. F. 2005 NewA, 11, 27
Condon, J. J., Cotton, W. D., Greisen, E. W. et al. 1998, AJ, 115, 1693
D'Abrusco, R., Longo, G., Walton, N. A. 2009 MNRAS , 396, 223
D'Abrusco, R., Massaro, F., Ajello, M. et al. 2012 ApJ, 748, 68
D'Abrusco, R., Massaro, F., Paggi, A. et al. 2013 ApJS, 206, 12
D'Abrusco, R., Massaro, F., Paggi, A. et al. 2014 ApJS, 215, 14
Di Mauro, M., Calore, F., Donato, F., Ajello, M., Latronico, L. 2014 ApJ, 780, 161
Falomo, R., Pian, E., Treves, A. 2014 A&ARv, 22, 73
Landoni, M., Massaro, F. et al. 2015 AJ, 149, 63
Massaro, E., Giommi, P., Leto, C. et al. 2009 A&A, 495, 691
Massaro, F., D'Abrusco, R., Ajello, M. et al. 2011a ApJ, 740L, 48
Massaro, F. & Ajello, M. 2011b ApJ, 729L, 12
Massaro, F., D'Abrusco, R., Tosti, G. et al. 2012a ApJ, 752, 61
Massaro, F., D'Abrusco, R., Tosti, G. et al. 2012b ApJ, 750, 138
Massaro, F., D'Abrusco, R., Giroletti, M. et al. 2013a ApJS, 207, 4
Massaro, F., D'Abrusco, R., Paggi, A. et al. 2013b ApJS, 206, 13
Massaro, F., D'Abrusco, R., Paggi, A. et al. 2013c ApJS, 209, 10
Massaro, F., Giroletti, M., Paggi, A. et al. 2013d ApJS, 208, 15
Massaro, F., Masetti, N., D'Abrusco, R. et al. 2014 AJ, 148, 66
Massaro, E., Maselli, A., Leto, C. et al. 2015b Ap&SS, 357, 75
Massaro, F., Landoni, M., D'Abrusco, R. et al.. 2015c, A&A, 575, 124
Massaro, F., D'Abrusco, R., Landoni, M. et al. 2015a ApJS, 217, 2
Massaro, F., Thompson, D. J., Ferrara, E. C. 2016 A&AR in press
Mattox, J. R., Schachter, J., Molnar, L., Hartman, R. C., Patnaik, A. R. 1997 ApJ, 481, 95
Mucke, A., Pohl, M., Reich, P., et al. 1997, A&A, 320, 33
Mahony, E. K., Sadler, E. M., Murphy, T., et al. 2010, ApJ, 718, 587
Mauch, T., Murphy, T., Buttery, H. J. et al. 2003 MNRAS, 342, 1117
Nolan, P. L., Abdo, A. A., Ackermann, M. et al. 2012 ApJS, 199, 31
Paggi, A., Massaro, F., D'Abrusco, R. et al. 2013 ApJS, 209, 9
Paggi, A., Milisavljevic, D., Masetti, N. et al. 2014 AJ, 147, 112
Ricci F., Massaro F., Landoni M. et al. 2015 AJ, 149, 160
Schinzel, F. K., Petrov, L., Taylor, G. B. 2015 ApJS, 217, 4
Shaw, M. S., Romani, R. W.; Cotter, G. et al. 2013 ApJ, 764, 135
Skrutskie, M. F. et al. 2006, AJ, 131, 1163
Stecker, F. W., Salamon, M. H. & Malkan, M. A. 1993, ApJ, 410, L71
Stickel, M., Padovani, P., Urry, C. M. et al. 1991 ApJ, 374, 431
Taylor, M. B. 2005, ASP Conf. Ser., 347, 29
Urry, C. M., & Padovani, P. 1995, PASP, 107, 803
Wright, E. L., et al. 2010 AJ, 140, 1868